AD)	

Award Number: DAMD17-99-1-9301

TITLE: Are p53 Mutations Associated with Increased Risk of Developing Breast Cancer? A Molecular Epidermiological Study

PRINCIPAL INVESTIGATOR: Rita A. Kandel, M.D.

CONTRACTING ORGANIZATION: Mount Sinai Hospital
Toronto, Ontario Canada M5G 1X5

REPORT DATE: October 2000

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release; Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

blank)	October 2000		p 99 - 1 Sep 00)		
4. TITLE AND SUBTITLE Are p53 Mutations Associ Developing Breast Cancer Study 6. AUTHOR(S)	5. FUNDING NUMBERS DAMD17-99-1-9301				
Rita A. Kandel, M.D.					
7. PERFORMING ORGANIZATION NAME Mount Sinai Hospita Toronto, Ontario Ca E-MAIL: rkandel@mtsinai.on.ca	1		8. PERFORMING REPORT NUI	G ORGANIZATION MBER	
9. SPONSORING / MONITORING AGE U.S. Army Medical Research and M Fort Detrick, Maryland 21702-501	10. SPONSORING / MONITORING AGENCY REPORT NUMBER				
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION / AVAILABILITY	ort contains colo	or photos		12b. DISTRIBUTION CODE	
Approved for public release; distrib	oution unlimited			125. DISTRIBUTION CODE	
detected by immunohistochemistr immunohistochemistry alone, we mutations in benign breast tissue at (1) analyzing benign breast tissue f DNA sequencing; and 2) estimating presence of both p53 mutations and with those detected in their preceed.	ested within a cohort of 4,888 y was associated with a 2.5-may have underestimated the associated with increased ristrom 138 cases and 556 controing the risk of breast cancer in d p53 protein accumulation in ling benign breast tissue samp.	-fold increase in the ri- e true risk of developings of subsequent breast of als for the presence of p5 in relation to: (a) the pro- BBD. We also propose les.	sk of subsequering breast cance cancer. We are to a mutations using esence of p53 metro compare mu	er. We hypothesized that p53 esting our hypothesis by: ang PCR-SSCP and PCR-direct nutations in BBD; and (b) the	

sections cut, epithelium enriched tissue microdissected out, and DNA extracted. The DNA has undergone PCR-SSCP under two conditions and those with abnormal gel patterns have undergone repeat PCR-SSCP. Abnormal migrating bands have been cut from the gel .These will be sequenced. We have determined that manual sequencing rather than automated sequencing is more appropriate to analyze the p53 gene in these small paraffin embedded benign lesions.

15. NUMBER OF PAGES 14. SUBJECT TERMS Breast Cancer 12 16. PRICE CODE 18. SECURITY CLASSIFICATION 19. SECURITY CLASSIFICATION 20. LIMITATION OF ABSTRACT 17. SECURITY CLASSIFICATION

OF REPORT Unclassified OF THIS PAGE Unclassified OF ABSTRACT Unclassified

Unlimited

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std. Z39-18

Table of Contents

Cover	1
SF 298	2
Table of Contents	3
Introduction	4
Body	4
Key Research Accomplishments	5
Reportable Outcomes	6
Conclusions	6
References	7
Appendices	8

INTRODUCTION

Our work is directed towards identifying genetic and protein changes in benign breast disease (BBD) which might be involved in the pathogenesis of breast cancer, and which might serve as markers of risk. We have recently completed a case-control study, nested within a cohort of 4,888 women with BBD, in which we demonstrated that p53 protein accumulation detected by immunohistochemistry was associated with a 2.5-fold increase in the risk of subsequent breast cancer (1). However, by using immunohistochemistry alone, we may have underestimated the true risk of developing breast cancer in association with p53 changes since approximately 33% of p53 mutations do not alter the protein in such a way that there is positive immunostaining (2-5). Therefore, a more complete assessment of the role of the association between p53 and breast cancer risk will come from studies combining both immunohistochemistry and p53 gene sequencing. We hypothesized that p53 mutations in benign breast tissue are associated with increased risk of subsequent breast cancer.

We are testing our hypothesis by:

- (1) analyzing benign breast tissue from 138 cases and 556 controls for the presence of p53 mutations using PCR-SSCP and PCR-direct DNA sequencing; and
- 2) estimating the risk of breast cancer in relation to: (a) the presence of p53 mutations in BBD; and (b) the presence of both p53 mutations and p53 protein accumulation in BBD. We will localize the mutations to determine whether they occur preferentially in specific sites of the DNA and to compare them to known mutations listed in p53 mutation banks (6-9). We also propose to compare mutations detected in the cancers with those detected in their preceeding benign breast tissue samples.

BODY

Task 1: Extracting DNA from paraffin blocks (Months 1-24):

- A) Cut histological sections from paraffin embedded tissue.
- B) Extract DNA

Task 2: PCR-SSCP analysis (Months 4-34):

- A) PCR for exons 2-11
- B) SSCP gels for each exon
- C) Autoradiography

We currently doing both task 1 and 2. To date we have generated primers for exons 2 to 11 of the p53 gene. We have developed the optimal PCR conditions, e.g. temperature, cycle number, primer concentration, and magnesium concentration, for each exon. We have cut histological sections from paraffin blocks (which contain the tissue to be analyzed), microdissected out the appropriate area in the tissue, and extracted the DNA using proteinase K from 152 subjects. We have examined the PCR products for each exon for 126 subjects. Each sample is run under two conditions (2 and 10% glycerol in

the loading buffer). Samples that show abnormal band migration in either one or both gels (please see a representative SSCP gel in the appendix 1) undergo repeat PCR. The new PCR product is then analyzed by repeat SSCP analysis. Those samples that show reproducible abnormal band migration are identified as samples that require DNA sequencing to confirm the presence of a p53 mutation or polymorphism. The bands are excised from the gels and stored at 4°C until they are further analyzed. By considering only those samples that show abnormalities in products from two separate PCRs as having gene alteration, we minimize the likelihood that the abnormality detected is an artefact resulting from nucleotide misincorporation during the PCR.

Task 3: Sequencing DNA with altered mobility on SSCP gels (Months 5-30):

- A) Excise band with altered mobility
- B) Elute DNA
- C) PCR appropriate exon(s)
- D) Sequence reaction and separation on acrylamide gels
- E) Autoradiography
- F) Automated sequencing

We are currently also working on task 3. Those samples that show reproducible abnormal band migration in the SSCP gels are identified as samples that require DNA sequencing to confirm the presence of a p53 mutation or polymorphism. The bands are excised from the SSCP gels and stored at 4°C until they are analyzed further. We are just beginning the next phase of eluting the samples and doing the sequencing.

We have compared the efficiency and reproducibility of automated sequencing to manual sequencing (part F). The Microgene Blaster TM (Visible Genetics, Toronto, CAN) which utilizes fluorescent labels was the automated sequencer used. This sequencer has been used successfully in other studies (10, 11). We obtained primers for exons 1 to 11 that were compatible with this sequencer. Using reagents obtained from Visible Genetics, sequencing of the PCR products was attempted. We found that if adequate amounts of DNA were used the p53 gene could be sequenced. However in our study we have only small amounts of DNA because unlike breast cancer, benign breast disease lesions are much smaller and thus we have much less starting tissue. We did detect base deletions and nucleotide changes in many of the samples examined but the alterations were not reproducible between analyses (see sequencing print out in appendix 2). All exons were examined to determine whether this problem was due to the method itself or specific to the exon and possibly the primers selected. All exons showed a similar lack of reproducibility. We concluded that for this study automated sequencing was not suitable. If possible we will examine whether the microarray technology might be more suitable.

KEY RESEARCH ACCOMPLISHMENTS

As we are just a year into the project for which the results are dependent on analysis of data from the entire case-control series, there are no key research accomplishments as of yet.

REPORTABLE OUTCOMES

There are no reportable outcomes as of yet.

Individuals who have been employed or paid from this grant include: Melissa Cooper MSc student ShuQiu Li Technician Tajinder Bhardwaj Technician (started July 2000) Hanje Chen Technician Shudong Zhang Post Doctoral Fellow (2 months)

CONCLUSIONS

We are able to extract DNA from the paraffin embedded tissue samples and the DNA obtained is suitable for PCR-SSCP and sequencing.

Gene abnormalities detected by PCR-SSCP will have to be confirmed using manual sequencing. Automated sequencing is not sufficiently sensitive or reproducible when using the small amounts of DNA extracted from the paraffin embedded tissue obtained in this study.

REFERENCES

- 1. Rohan TE, Hartwick W, Miller AB, Kandel R. Immunohistochemical-detection of c-erbB-2 and p53 in benign breast disease and breast cancer risk. J Nat'l Cancer Institute 1998;90:1262-1269.
- 2. Dunn JM, Hastrich DJ, Newcomb P, Webb JCJ, Maitland NJ, Farndon JR. Correlation between p53 mutations and antibody staining in breast carcinoma. Br J Surg 1993;80:1410-12.
- 3. Jacquemier J, Molès JP, Penault-Llorca F, Adélaide J, Torrente M, Vineis P, Birnbaum D, Theillet C. p53 immunohistochemical analysis in breast cancer with four monoclonal antibodies: comparison of staining and PCR-SSCP results. Br J Cancer 1994;69:846-52.
- 4. Sjögren S, Inganäs M, Norberg T, Lindgren A, Nordgren H, Holmberg L, Bergh J. The p53 gene in breast cancer: Prognostic value of complementary DNA sequencing versus immunohistochemistry. . J Nat'l Cancer Institute 1996;88:173-82.
- 5. Visscher DW, Sarkar FH, Shimoyama RK, Crissman JD. Correlation between p53 immuno- staining patterns and gene sequence mutations in breast carcinoma. Diagn Mol Pathol 1996; 5:187-93.
- 6. Beroud C, Verdier F, Soussi T. p53 gene mutation: software and database. Nucleic Acids Res 1996; 24:147-50.
- 7. De Vries EMG, Ricke DO, De Vries TN, Hartmann A, Blaszyk H, Liao D, Sousi T, Kovach JS, Sommer SS. Database of mutations in the p53 and APC tumor suppressor genes designed to facilitate molecular epidemiological analyses. Hum Mut 1996;7:202-13.
- 8. Hainaut P, Soussi T, Shomer B, Hollstein M, Greenblatt M, Hovig E, Harris CC, Montesano R. Database of p53 gene somatic mutations in human tumors and cell lines: updated compilation and future prospects. Nucleic Acids Res 1997;25:151-7.
- 9. Hollstein M, Shomer B, Greenblatt M, Soussi T, Hovig E, Montesano R, Harris CC. Somatic point mutations in the p53 gene of human tumors and cell lines: updated compilation. Nucleic Acids Res 1996;24:141-6.
- 10. Bharaj BS, Angelopoulou K, Diamandis EP. Rapid sequencing of the p53 gene with a new automated sequencer. Clin Chem 1998;44:1397-1403.
- 11. Saunders ME, MacKenzie R, Shipman R, Fransen E, Gilbert R, Jordan RC.Patterns of p53 gene mutations in head and neck cancer: full-length gene sequencing and results of primary radiotherapy. Clin Cancer Res. 1999;5:2455-63.


APPENDIX 1

PCR-SSCP GEL SHOWING 33 SAMPLES.

Altered mobility is seen in samples. The bands will be excised and the DNA sequenced to determine whether the sequence change represents polymorphism(s) or mutation(s). No PCR product was obtained for sample #92.

PCR-SSCP p53 EXON 3

31/53/55/57/58/65/70/72/79/83/92/124/126/175/187/188/189/190/191/192/195/196/197/199/201/203/205/206/207/208/220/222/224



APPENDIX 2

Sequences of exon 4 of the p53 gene obtained from DNA extracted from paraffin embedded breast tissue showing benign breast disease changes. The sample was run on two separate occasions (A,B) and different sequences were obtained.

lab200001261141,1-4 - p53 Breast Tumors

Machine: 701098

son!

Date: 20000126

Operator: Bhupinder

5 Ö 800 CTGG 1GC CT GGCKTC [2] 280 ပ O B B O

306

305

298

1286 1290 1294

18 | 282

11 | 278

<u>g</u>,

S G G C C

15) 15)

File: /Users/lab/Data/p53 Breast Tumors/Assays/Jan2000/Bhupinder/lab200001261141,1-4.gel

116 159 161 163 A10 1312 168 E 210 155 157 159 160 161 162, 164 166 1304 1306 1308 **≻** O 109 111 157 **[251 | 253 | 255** ပ ပ 204 |299 | |302 | 205 247 |249 150 1103 1105 G C C ပြ O 199 1778 1180 1182 1184 1186 1188 1190 1192 1194 1196 245 -967 -236-| |146 -101 F |51 | |53 243 Ω |291 |293 |294 142 1144 98 S S G --96 -64 238 239 10 1134 1136 1138 1140 94 22255 288 92 124 1232 1234 1236 133' 135' 10 138' 140' 142' 144 lo 188 lo 190 |5<mark>4</mark>- |0| C C ົວ Ö 1269 1271 1273 1275 1278 ැ **ග** . 28 228 129 8 1,64 1,66 1,68 1,70 1,72 1,74 1,76 223 225 Date: 20000118
Assay Notes: Pc R 13.1.200
GeneObjects 3.1 (1999/07/13 20:28:45 EDT) |78 |79 | **|81** O G C 219 221 1267 (ග 76 -82 1216 263 | 265 -82 -129 74 |23 |25 | _0 116 118 ් 212 g 696 200 0 2677

036

280

Operator: Bhupinder

lab200001180945,1-4 - p53 Breast Tumors

Machine: 701098

File: /Users/lab/Data/p53 Breast Tumors/Assays/Jan2000/Bhupinder/lab200001180945,1-4.gel

Table of Contents

Cover	1
SF 298	2
Table of Contents	3
Introduction	4
Body	4
Key Research Accomplishments	5
Reportable Outcomes	6
Conclusions	6
References	7
Appendices	8